

AD-A238 594

2

SECURITY CLASSIFICATION OF THIS PAGE



REPORT DOCUMENTATION PAGE

OMB No. 0704-0188

1a REPORT SECURITY CLASSIFICATION (u)			1b RESTRICTIVE MARKINGS NA		
2a SECURITY CLASSIFICATION AUTHORITY NA			3 DISTRIBUTION/AVAILABILITY OF REPORT Distribution Unlimited		
2b DECLASSIFICATION/DOWNGRADING SCHEDULE NA					
4 PERFORMING ORGANIZATION REPORT NUMBER(S) Duke University Medical Center			5 MONITORING ORGANIZATION REPORT NUMBER(S) NA		
6a NAME OF PERFORMING ORGANIZATION Duke University Medical Center		6b OFFICE SYMBOL (if applicable) NA	7a NAME OF MONITORING ORGANIZATION Office of Naval Research		
6c ADDRESS (City, State, and ZIP Code) Duke University Medical Center Box 3181 Durham, NC 27710			7b ADDRESS (City, State, and ZIP Code) 800 N. Quincy Street Arlington, VA 22217-5000		
8a NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research		8b OFFICE SYMBOL (if applicable) ONR	9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N00014-88-k-0132/P00001		
8c ADDRESS (City, State, and ZIP Code) 800 N. Quincy Street Arlington, VA 22217-5000			10 SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO 61153N	PROJECT NO RR04108	TASK NO 4414804
			WORK UNIT ACCESSION NO		
11 TITLE (Include Security Classification) Models of Excitation-Secretion Coupling in Pituitary Cells					
12 PERSONAL AUTHOR(S) C. Frank Starmer, Ph.D.					
13a TYPE OF REPORT Final		13b TIME COVERED FROM 11/89 TO 10/90		14 DATE OF REPORT (Year, Month, Day) 6/26/91	
				15 PAGE COUNT 14	
16 SUPPLEMENTARY NOTATION					
17 COSATI CODES			18 SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD GROUP SUB-GROUP			mathematical model, ion channel, GH3 cell, neural model		
19 ABSTRACT (Continue on reverse if necessary and identify by block number) <p>This report describes final studies in developing a biophysical description of the electrical events surrounding hormone release in pituitary cells. Utilizing a model based on a membrane capacitance shunted by potassium and calcium channels, we are developing a computer program that will simulate the dynamic response of the transmembrane potential of GH3 cells in response to thyrotropin-releasing hormone (TRH). When available, we are using published channel models of potassium and calcium channels. In parallel, we are pursuing a phase-plane description of the electrical properties. These models and analyses will be used to investigate the effect of temperature on membrane action potentials.</p>					
20 DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS			21 ABSTRACT SEC (u)		
22a NAME OF RESPONSIBLE INDIVIDUAL Dr. J.A. Majde			22b TELEPHONE (Include Area Code) (202) 696-4055		22c OFFICE SYMBOL ONR

91-03873



**Best
Available
Copy**

Research Contract N00014-88-K-0132
Final Report

Principal Investigator: C. Frank Starmer, Ph.D.
Duke University Medical Center

NT'S GRAAL <input checked="" type="checkbox"/>	
DTIC TAB <input type="checkbox"/>	
Unannounced <input type="checkbox"/>	
Justification <input type="checkbox"/>	
By _____	
Distribution/ _____	
Availability Codes _____	
Dist	Avail and/or Special
A-1	

Introduction:

Hormone release from pituitary cells appears related to a modification of the basal electrical activity of the cell. This electrical activity has been shown to be modulated by agents that also regulate secreting activity. For instance Thyrotropin-releasing hormone (TRH) triggers the release of prolactin in GH₃ cells and simultaneously leads to an increase in action potential frequency in these electrically active cells. The membrane potential associated with the appearance of released hormone in the extracellular fluid appears to initially be hyperpolarized (thought to result in opening of a Ca⁺ activated potassium channel) followed by a decrease in the voltage dependent K⁺ currents. Dubinsky and Oxford (1) have suggested that upon application of TRH 1: CA⁺⁺ is released from intracellular stores which activates CA⁺⁺ activated K channels; 2: voltage-dependent K channel openings are depressed during hyperexcitable phase and that 3: TRH does not directly modulate calcium channel activity. During the burst of action potentials during the hyperexcitable phase, extracellular CA⁺⁺ enters the cells through voltage gated CA⁺⁺ channels (perhaps to participate in prolactin secretion) (2) and accumulates to a point where electrical activity becomes again silent.

Temperature effects the transition rates of voltage gated channels (3,4). This project addresses the role of channel gating on models of the electrical properties of cellular action potentials and on the accumulation of intracellular Ca⁺⁺. The work has followed three directions: that of developing computing tools to facilitate management and display of model results; developing a minimal model of the electrical properties of GH₃ cells and to develop analytic tools to characterize the use-and frequency-dependent properties of the accumulation intracellular calcium during bursts of action potentials.

Progress Report:

1) Our research group has been involved in developing software tools to facilitate acquisition, management, analysis and display of research data. We have taken the approach that all data files must be self-describing, e.g. files that control experiments such as stimulus protocols as well as primary data files must be readily interpretable without the use of other documents. To accomplish this goal, we include in self-documenting files a dictionary record (as the first record of the file) that names the variables stored in the file. This record is followed by one or more "comment" records that describe the experiment and the experimental conditions. Together with the raw data, these records define a file that can be easily interpreted.

The most recent work has focused on developing tools for visualizing primary data, results of simulations and displays of derived results. Using the X-window system operating on Sun 4 workstations, we have developed a graphical editor that allows cutting and pasting segments of a graphic display. These selected segments can be printed or subjected to further analyses e.g. curve fitting. In addition we have developed a tool for scanning a sequence of research data records (e.g. simulations for a set of different conditions). These tools have been extremely useful in visually investigating a small segment of a long simulation. The software tools have been described in several manuscripts listed at the end of this report (7,8).

2) GH₃ cell model. For a minimal model, we have considered a 3 component model: a voltage activated potassium current, I_K, and voltage and calcium inactivated calcium current, I_{Ca}, and a leakage current, I_l, that are considered electrically in parallel with the membrane capacitance. In addition, we have assumed a first order sequestration of internal calcium. Defining the membrane capacitance as C_m we define the minimal model as

$$\frac{dv}{dt} = - \frac{1}{C_m} (I_{Ca} + I_K + I_l)$$


where V is the membrane potential, I_{Ca} is the calcium current described by

$$I_{Ca} = \bar{g}_{Ca} d(V) f(V, Ca_i) (V - V_{Ca})$$

where d and f are activation and inactivation gating variables, V_{Ca} is the calcium reversal potential and \bar{g}_{Ca} is the maximum calcium conductance. Similarly, for the potassium current

$$I_K = \bar{g}_K n(V) (V - V_K)$$

where $n(V)$ is the activation variable, V_K is the reversal potential and \bar{g}_K is the maximum potassium current. The gating variables are defined to reflect transitions in channel protein conformations according to a simple first order model



so that for the potassium channel, n at equilibrium is $n = \alpha / (\alpha + \beta)$ and $n(t)$ is the solution to

$$\frac{dn}{dt} = \alpha (1 - n) - \beta n$$

To incorporate Ca^{++} inactivation into the inactivation variable f , we assume a first order process leading to an equilibrium inactivation of the form $f_{\infty} = [1 + Ca^{++}/k_n]^{-1}$. Finally, intracellular calcium distribution is determined by

$$\frac{d}{dt} Ca^{++} = I_{Ca} - K_{Ca} Ca_i^{++}$$

where I_{Ca} represents calcium entering the cell through open channels and is normalized per unit volume. We have solved these equations with some preliminary estimates of rate constants and have investigated temperature dependence assuming a Q_{10} of 3. We have found that this simple model

can exhibit 3 types of temperature sensitive behavior (figure 1 and 2): simple oscillations, bursts of oscillations and continuous oscillations from a depolarized baseline. These results suggest the nonlinear terms in the model can produce behavior similar to that seen in other systems exhibiting behavior described by the term, chaos. Figure 1 illustrates results when the absorption rate of Ca_i is held constant while the temperature effect is restricted to channel conformation rates. Figure 2 illustrates the same sets of rate constants but also allowing k_{Ca} to vary with temperature. The vertical axis is membrane potential (mV) while the horizontal axis is time (msec). The detailed mechanism leading to such dramatic changes in oscillatory behavior is the focus of current investigations.

3) Our work with analytically characterizing the use-dependent properties of Ca_i has followed that of our models of ion channel blockade (5). Basically, with each action potential, intracellular calcium is incremented by a fraction, proportional to the difference between intracellular and extracellular calcium. For the n^{th} action potential when the channel is conducting

$$\frac{d}{dt}C_i = \gamma(C_o - C_i) - K_a C_i$$

where C_i and C_o are intracellular and extracellular concentrations and γ represents diffusion rate down the calcium concentration gradient. When the channel is not conducting, C_i is reduced through intracellular storage at a rate K_a so

$$\frac{d}{dt}C_i = -K_a C_i$$

If the channel open time is exponentially distributed with mean, t_o , (5) then

$$C_i = C(\infty) + [C(0) - C(\infty)] e^{-(\gamma C_o + k_a)t_o}$$

where $C(\infty) = \gamma C_0 / (\gamma + k_a)$

While the channel is not conducting

$$C_i(t) = C(o)e^{-k_a t}$$

During a burst, internal calcium is incrementally increased so that for the n^{th} pulse

$$C_n = C_{ss} + (C_0 - C_{ss})e^{-[k_a t_r + (\gamma + k_a)t_0]n}$$

where t_r is the interpulse interval between action potentials during a burst and t_0 is the mean channel open time and

$$C_{ss} = \frac{C(\infty)(1 - e^{-(\gamma + k_a)t_0}}{1 - e^{-[k_r t_r + (\gamma + k_a)t_0]}}$$

Thus, it is possible to estimate the behavior of intracellular calcium during and between bursts if the channel open time is exponentially distributed. These preliminary analyses indicate that temperature influences on channel gating will modify the oscillatory bursting properties of the cells.

4) The role of channel gating in drug-complexed channels. We have explored the role of pharmacologic blockade of ion channels using data from our own laboratory as well as data from studies of the interaction of nimodipine with calcium channels from GH₄C₃ cells (6). Measurements of ionic currents are described by Ohms law as

$$I_{Ca} = g_{Ca} d f (1 - b) (V - V_{Ca})$$

where d and f are activation and inactivation parameters determined from the channel gating properties, b is the fraction of blocked channels and V_{Ca} is the calcium reversal potential. We found that it is not possible to estimate drug binding properties from measurements of ionic currents if the drug both blocks the channel and modifies the activation and/or inactivation process. The reason is that blockade,

activation and inactivation may share the same voltage dependence. Unless binding is via a fixed affinity site, it is not possible to extract unique and unambiguous voltage dependent rates. For example, any change in the voltage dependence of channel gating could be negated by a postulated voltage dependence of drug binding. We are the first to demonstrate that it is not possible to uniquely identify channel gating parameters in the presence of variable affinity binding. These results, describing both studies of lidocaine blockade of sodium channels and nimodipine block of calcium channels were recently published(10 in Contract Related Publications).

References


1. Dubinsky, J.M. and Oxford, G.S. Dual modulation of K channels by thyrotropin-releasing hormone in clonal pituitary cells. Proc. Natl. Acad. Sci. 82:4282-4286, 1986.
2. Benham, C.D. Voltage-gated and agonist-mediated rises in intracellular Ca^{++} in rat clonal pituitary cells (GH₃) held under voltage clamp. J. Physiol. (London) 415:143-158, 1989.
3. Hodgkin, A.L. and Huxley, A.F. A quantitative description of membrane current and its application to conduction and excitation in nerve. J. Physiol. (London) 117:500-544, 1952.
4. Partridge, L.D. and Conner, J.A. A mechanism for minimizing temperature effects on repetitive firing frequency. Amer. J. Physiol. 234:C155-C161, 1978.
5. Starmer, C.F. Characterizing activity dependent processes with a piecewise exponential model. Biometrics, 44:549-559, 1988.
6. Cohen, C.J. and McCarthy, R.T. Nimodipine block of calcium channels in rat anterior pituitary cells. J. Physiol (London) 387:195-225, 1987.

Contract Related Publications


1. Starmer, CF, Nesterenko, VV, Undrovinas, A.I., Packer, D.L., Gilliam, F.R., Grant, A.O., Rosenshtraukh, L.V. and Strauss, H.C. Characterizing ion channel blockade with the guarded receptor hypothesis. Molecular and Cellular Mechanisms of Antiarrhythmic Agents. Futura, Mt Kisco, NY. 1988
2. Starmer, C.F. Characterizing activity dependent processes with a piecewise exponential model. Biometric 44:549-559, 1988.
3. Starmer, C.F., Undrovinas, A.I., Scamps, .F, Vassort, G., Nesterenko, V.V., and Rosenshtraukh, L.V. Ethacizin blockade of Ca^{++} channels: A test of the guarded receptor hypothesis. Amer. J. Physiol. 257:H1693-H1704, 1989.
4. Gilliam, F.R., Starmer, C.F. and Grant, A.O. Blockade of rabbit atrial sodium channels by lidocaine: Characterization of continuous and frequency dependent blocking. Circ. Res. 65:723-739, 1989.
5. Packer, D.L., Grant, A.O., Strauss, H.C. and Starmer, C.F. Characterization of concentration and use-dependent effects of quinidine from conduction delay and declining conduction velocity in canine Purkinje fibers. J. Clin. Invest. 83:2109-2119, 1989.
6. Grant, A.O., Dietz, M.A., Gilliam, F.R. and Starmer, C.F. Blockade of cardiac sodium channels by lidocaine and its hydrophobic derivative RAD-242: Single channel analysis. Circ. Res. 65:1247-1262, 1989.
7. Starmer, C.F. and Dietz, M.A. Managing clinical research data: Software tools for hypothesis exploration. Environmental Health Perspectives. 87:5-11, 1990.
8. Dietz, M.A., Grant, A.O. and Starmer, C.F. An object

oriented user interface for analysis of biological data.
Computers and Biomedical Res. 23:82-96, 1990.


9. Whitcomb, D.C. Gilliam, F.R., Starmer, C.F. and Grant, A.O. Marked QRS complex abnormalities and sodium channel blockade by propoxyphene reversed with lidocaine. J. Clin. Invest. 84:1629-1636, 1989.
10. Starmer, C.F., Nesterenko, V.V., Gilliam, F.R. and Grant, A.O. Using ionic currents to identify and estimate parameters in models of channel blockade. Amer. J. Physiol 259:H626-H634, 1990.
11. Gilliam, F.R., Rivas, P.A., Wendt, D.J., Starmer, C.F. and Grant, A.O. Extracellular pH modulates the block of both sodium and calcium channels by Nicardipine. Am. J. Physiol. 259:H1178-H1184, 1990.
12. Barber, J.M., Starmer, C.F. and Grant, A.O. Evidence of two use-dependent binding sites in sodium channels of isolated rabbit myocytes. Circ. Res. (In press).



$T=25^{\circ}\text{C}$, $K_{\text{ca}}=.04/\text{msec}$



$T=20^{\circ}\text{C}$, $K_{\text{ca}}=.04/\text{msec}$



$T=17^{\circ}\text{C}$, $K_{\text{ca}}=.04/\text{msec}$

Figure 1






Figure 2

DISTRIBUTION LIST

Stress Neurochemistry Program

Annual, Final and Technical Reports (one copy each)

INVESTIGATORS

Dr. H. Elliott Albers
Lab. Neuroendocrin. & Behavior
Depts. of Biology & Psychology
Georgia State University
Atlanta, GA 30303

Dr. Gwen V. Childs
Dept. of Anatomy & Neuroscience
Univ. of Texas Medical Branch
Galveston, TX 77550

Dr. Carl E. Creutz
Dept. of Pharmacology
University of Virginia
Charlottesville, VA 22908

Dr. Mary F. Dallman
Dept. of Physiology
University of California, Box 0444
San Francisco, CA 94143-0444

Dr. Caleb E. Finch
Dept. of Neurobiology
Univ. of Southern California
Los Angeles, CA 90089-0191

Dr. Thackery S. Gray
Department of Anatomy
Loyola University Medical Center
216 South First Avenue
Maywood, IL 60153

Dr. Richard F. Ochillo
College of Pharmacy
Xavier Univ. of Louisiana
7325 Palmetto Street
New Orleans, LA 70125

Dr. Terry Reisine
Dept. of Pharmacology
Univ. of Pennsylvania
School of Medicine
36th and Hamilton Walk
Philadelphia, PA 19104

Dr. C. Frank Starmer
P.O. Box 3181
Duke Univ. Medical Center
Durham, NC 27710

Dr. Kent E. Vrana
Dept. of Biochemistry
West Virginia School
of Medicine
Morgantown, WV 26506

Stress Neurochemistry

Annual, Final and Technical Reports (one copy each except as noted)

ADMINISTRATORS

Scientific Officer, Physiology
Code 1141S8
Office of Naval Research
800 N. Quincy Street
Arlington, VA 22217-5000

Program Manager, Code 1213
Human Factors Biosciences
Division
Office of Naval Research
800 N. Quincy Street
Arlington, VA 22217-5000

Administrator (2 copies) (Enclose DTIC Form 50)
Defense Technical Information Center
Building 5, Cameron Station
Alexandria, VA 22314

Program Manager, Code 223
Support Technology
Directorate
Office of Naval Technology
800 N. Quincy Street
Arlington, VA 22217-5000

Administrative Contracting Officer
ONR Resident Representative
(address varies - obtain from business office)

Annual and Final Reports Only (one copy each)

DoD ACTIVITIES

Commanding Officer
Naval Medical Center
Washington, DC 20372

Commanding Officer, Code 404
Naval Medical Research & Development Command
National Naval Medical Center
Bethesda, MD 20814

Commander
Chemical and Biological Sciences Division
Army Research Office, P.O. Box 12211
Research Triangle Park, NC 27709

Directorate of Life Sciences
Air Force Office of Scientific Research
Bolling Air Force Base
Washington, DC 20332

Final and Technical Reports Only

Director, Naval Research Laboratory (6 copies)
Attn: Technical Information Division, Code 2627
Washington, DC 20375